

A Reference Guide To REIMBURSEMENT & CODING

FOR ALL BILLING OR CODING INQUIRIES
Call 1-800-473-1430 or E-mail: Cipla.Lanreotide@cipla.com

Lanreotide Injection

(lanreotide acetate) for injection

120 mg

FDA APPROVED: DECEMBER 17, 2021¹

INDICATIONS

Lanreotide Injection is indicated for the treatment of adult patients with unresectable, well or moderately differentiated, locally advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NETs) to improve progression-free survival.

IMPORTANT SAFETY INFORMATION

Contraindications: Hypersensitivity to lanreotide.

Warnings and Precautions

- **Cholelithiasis and Complications of Cholelithiasis:** Monitor patients periodically. Discontinue if complications of cholelithiasis are suspected. Gallstones may occur; consider periodic monitoring.
- **Hyperglycemia and Hypoglycemia:** Glucose monitoring is recommended and antidiabetic treatment adjusted accordingly.

[Please see Page 10 for additional Important Safety Information.](#)



The information provided herein is for information and reference purposes only. It is the responsibility of the provider to determine coverage, reimbursement and appropriate coding for a particular patient, product and/or procedure. The information herein is not a guarantee of coverage or reimbursement for Lanreotide or any other product or services.

[Click here for Full Prescribing Information.](#)

www.ciplalanreotide.com

Medical Information:

Report Adverse Events/Side Effects
Report Product Complaints
Submit Medical Inquiries:

Contact Us:  1-866-604-3268  DrugSafety@cipla.com

Cipla USA, Inc.
10 Independence Blvd., Suite 300
Warren, NJ 07059

Customer Service: 844-CIPLA US (844-247-5287)

Email: Ciplacs@ups.com

TABLE OF CONTENTS

INTRODUCTION 3

HCPCS CODES 4

PRODUCT PRICING 4

PRODUCT SPECIFIC HCPCS CODES,
 PHYSICIAN OFFICE AND HOPD 4

NOT OTHERWISE CLASSIFIED (NOC) HCPCS CODES 4

PRODUCT SPECIFIC HCPCS CODES,
 PRODUCT PRICING 4

MEDICARE PAYMENT RATES: PHYSICIAN OFFICE (MPFS)/
 HOSPITAL OUTPATIENT (OPPS) 4

DIAGNOSIS CODES 5

ICD-10-CM DIAGNOSIS CODES FOR ACROMEGALY 5

ICD-10-CM DIAGNOSIS CODES FOR GEP-NETS 5

CURRENT PROCEDURAL TERMINOLOGY (CPT) 6

DRUG ADMINISTRATION/CPT CODES 6

NATIONAL DRUG CODES 7

PRODUCT INFORMATION FOR LANREOTIDE 7

PHYSICIAN OFFICE 8

SAMPLE CMS-1500 CLAIM FORM 8

HOSPITAL OUTPATIENT 9

SAMPLE CMS-1450 CLAIM FORM 9

REFERENCES 10

FULL PRESCRIBING INFORMATION 11

[Click here for Full Prescribing Information.](#)

www.ciplalanreotide.com

At Cipla, We Are Committed to Providing Helpful Reimbursement Information for Physician Offices and Facilities to Ensure Patient Access to Lifesaving Medicines

Invagen Pharmaceuticals, Inc., an affiliate of Cipla USA, Inc. ("Cipla") received FDA final approval for its LANREOTIDE Injection supplied in 120 mg/0.5 mL single-dose pre-filled syringes ("LANREOTIDE") on December 17, 2021, under a New Drug Application ("NDA") submitted under the 505(b)(2) filing pathway.¹ The active ingredient, the strength (120 mg) and dosage (once every four weeks via deep subcutaneous injection, the same as Somatuline Depot[®] from Ipsen Biopharmaceuticals Inc.,) for the treatment of gastroenteropancreatic neuroendocrine tumors (GEP-NETs).¹

Cipla has applied to the Centers for Medicare & Medicaid Services (CMS) and based on CMS' HCPCS Decision Tree, Cipla believes that J1930 is still applicable for Cipla's Lanreotide 120 mg.²⁹

The information provided in this guide is for informational and reference purposes only. Coding and coverage policies can change, often without warning. It is the provider's responsibility to determine coverage and appropriate coding for a particular patient and/or procedure. The information in this guide is not a guarantee of coverage or reimbursement for any product or service. Please contact your patient's health plan to confirm coding for a specific plan.

Until a permanent J-code is assigned to LANREOTIDE, it is common practice for payers to require one or more miscellaneous HCPCS codes for coding and billing. When deciding which code best describes the services provided, refer to the payer's specific policy (i.e., Medicaid, Commercial Payer) for guidance on the appropriate miscellaneous HCPCS code(s) for LANREOTIDE. For Medicare Part B, please refer to the local Medicare Administrative Contractor (MAC) jurisdiction for guidance on HCPCS code requirements for unclassified drugs until CMS assigns a permanent code.⁴

[Click here for Full Prescribing Information.](#)

www.ciplalanreotide.com

PRODUCT SPECIFIC HCPCS CODES, PHYSICIAN OFFICE and HOPD⁴

J1930*	Injection, Lanreotide, 1 mg	Billing Unit: 1 mg
--------	-----------------------------	--------------------

*Cipla has applied to the Centers for Medicare & Medicaid Services (CMS) and based on CMS' HCPCS Decision Tree, Cipla believes that J1930 is still applicable for Cipla's Lanreotide 120 mg.²⁹

NOT OTHERWISE CLASSIFIED (NOC) HCPCS CODES⁴

Physician Office or Hospital Outpatient	<p>NOC Billing Unit: 1</p> <p>Include drug-identifying information in the comment field of the claim form, as specified by the payer</p>
J3490 (unclassified drugs)	
J9999 (not otherwise classified, anti-neoplastic drugs)	
Hospital Outpatient Who Bills Under the Outpatient Prospective System (OPPS), Typically Used for Medicare	
C9399 (unclassified drugs or biologicals)	

The payment allowance limits for drugs and biologicals that are not included in the ASP Medicare Part B Drug Pricing File or Not Otherwise Classified (NOC) Pricing File, are based on 103% of the WAC, or invoice pricing if the WAC is not published, except under OPPS where the payment allowance limit is AWP-5%.^{5,6}

PRODUCT PRICING^{7,8}

Wholesale Acquisition Cost (WAC)	\$8892.66 per 120mg/0.5 ml pre-filled syringe
Average Wholesale Price (AWP)	\$10671.19 per 120mg/0.5 ml pre-filled syringe

NOC MEDICARE PAYMENT RATES: PHYSICIAN OFFICE (MPFS) / HOSPITAL OUTPATIENT (OPPS)⁷

MPFS, NOC Payment Rate (No ASP established)	WAC + 3%
OPPS, NOC Payment Rate (No ASP established)	AWP – 5%

[Click here for Full Prescribing Information.](#)

Diagnosis Codes

The following may be appropriate ICD-10-CM diagnosis code for LANREOTIDE (This list of ICD-10-CM codes is not exhaustive. Providers should report the appropriate diagnosis code for each patient's condition.):

ICD-10-CM DIAGNOSIS CODES FOR ACROMEGALY⁹

Acromegaly and pituitary gigantism	E22.0
---	-------

ICD-10-CM DIAGNOSIS CODES FOR GEP-NETS⁹

Malignant carcinoid tumor of the duodenum	C7A.010
Malignant carcinoid tumor of the jejunum	C7A.011
Malignant carcinoid tumor of the ileum	C7A.012
Malignant carcinoid tumor of the small intestine, unspecified portion	C7A.019
Malignant carcinoid tumor of the appendix	C7A.020
Malignant carcinoid tumor of the cecum	C7A.021
Malignant carcinoid tumor of the transverse colon	C7A.023
Malignant carcinoid tumor of the descending colon	C7A.024
Malignant carcinoid tumor of the sigmoid colon	C7A.025
Malignant carcinoid tumor of the rectum	C7A.026
Malignant carcinoid tumor of the large intestine, unspecified portion	C7A.029
Malignant carcinoid tumor of the stomach	C7A.092
Malignant carcinoid tumor of the foregut, unspecified	C7A.094
Malignant carcinoid tumor of the midgut, unspecified	C7A.095
Malignant carcinoid tumor of the hindgut, unspecified	C7A.096
Malignant poorly differentiated neuroendocrine tumors	C7A.1
Other malignant neuroendocrine tumors	C7A.8
Secondary carcinoid tumors, unspecified site	C7B.00
Secondary carcinoid tumors of distant lymph nodes	C7B.01
Secondary carcinoid tumors of peritoneum	C7B.04
Secondary carcinoid tumors of other sites	C7B.09
Other secondary neuroendocrine tumors	C7B.8
Malignant neoplasm of ampulla of Vater	C24.1
Malignant neoplasm of endocrine pancreas	C25.4

Additional ICD-10-CM diagnosis codes may be used to fully describe the patient's condition and associated manifestations.

[Click here for Full Prescribing Information.](#)

Current Procedural Terminology (CPT) Drug Administration Codes*

The following CPT® code may be appropriate to report LANREOTIDE administration services. Evaluation and Management (E&M) codes for office visit services in addition to injection may be appropriate. Most payers require documentation of a separate and identifiable procedure. Some payers may not allow for a level 1 office visit and an injection code to be billed for the same date of service and may only allow for other levels of office visits to be billed with an appropriate modifier.

The CPT codes that may be appropriate for administration of LANREOTIDE appear in the table below.¹⁰

DRUG ADMINISTRATION/CPT CODES ¹⁰	
Therapeutic, prophylactic, or diagnosis injection; subcutaneous or intramuscular	96372
Chemotherapy administration, subcutaneous or intramuscular; hormonal anti-neoplastic	96402

* CPT codes and descriptions only are ©2019 by American Medical Association (AMA). All rights reserved. The AMA assumes no liability for data contained or not contained herein. CPT is a registered trademark of the American Medical Association.

[Click here for Full Prescribing Information.](#)

www.ciplalanreotide.com

National Drug Codes (NDCs)

Drug products are identified and reported using a unique, three-segment number, called the National Drug Code, a universal product identifier. The NDC is used primarily for pharmacy claims. However, it may also be required when billing physician-administered drugs to ensure crosswalk accuracy. When providers must include an NDC on an insurance claim, it typically must be in the required 11-digit format.

PRODUCT INFORMATION FOR LANREOTIDE¹

NDC: Payer requirements regarding the use of a 10-digit or 11-digit NDC vary

	10-DIGIT	11-DIGIT
LANREOTIDE ACETATE (120 mg/0.5mL)	69097-870-67	69097-0870-67
Dosage forms and strengths	GEP-NETs: 120 mg every 4 weeks.	
Route of Administration	For deep subcutaneous injection only	

[Click here for Full Prescribing Information.](#)

www.ciplalanreotide.com

Physician Office: Sample CMS-1500 Claim Form¹¹

Lanreotide and the associated services provided in the physician's office are billed on the CMS-1500 claim form or its electronic equivalent (837P)

Note: Sample form has been provided for informational purposes only. Providers should contact their patient's health plans for specific information on their coding requirements¹¹

19

Product Information –
When using miscellaneous code J3490 to report the use of LANREOTIDE, include the NDC number (e.g., 69097-0870-67), name of the drug, strength of the drug administered, and dose¹²

21

Diagnosis Code –
Enter appropriate ICD-10-CM diagnosis code(s) (e.g., C7A.011 - Malignant carcinoid tumor of the jejunum) Code to the highest specificity supported by the medical record¹³

24A

Dates of Service & NDC –
Enter the date(s) of service, and in the shaded area above the line, the required NDC for which a drug is reported in 24D. The NDC is preceded by the qualifier N4 and followed by the quantity qualifier (UN) and the quantity administered beginning in position¹⁴

24D

Drug Code –
Enter the appropriate HCPCS drug and CPT administration codes.
For Example: Drug – J3490 (unclassified drug), **Administration –** 96402 (Chemotherapy administration, subcutaneous or intramuscular; hormonal anti-neoplastic)¹⁵

24E

Diagnosis Code Pointer –
Specify the diagnosis from Box 21 that relates to the drug or procedure in¹⁶

24G

Days or Units –
Enter the number of service units for each line item. For a miscellaneous code, bill 1 unit¹⁷

Note: When billing for unlisted drugs using a NOC code, field 19 (or the equivalent electronic Loop/Segment in the 837-P, must be populated with the specific name of the drug(s) and/or a short descriptor of the drug(s)¹⁸

Click here for Full Prescribing Information.

Hospital Outpatient: Sample CMS-1450 Claim Form¹⁹

Lanreotide and the associated services provided in the hospital outpatient clinic or office are billed on the CMS-1450 claim form or its electronic equivalent (837I).²⁰

Note: Sample form has been provided for informational purposes only. Providers should contact their patient's health plans for specific information on their coding requirements¹⁹

The form contains the following key information:

- FL42 Revenue Code:** 0636 LANREOTIDE (lanreotide acetate) 120mg 1 unit, 0331 Chemo admin, injected
- FL43 Revenue Code Description:** J3490 (Drug), 96402 (Chemo admin)
- FL44 Drug Code:** J3490 (Drug), 96402 (Administration)
- FL45 Date of Service:** 10-01-22
- FL46 Service Units:** 1
- FL66 Diagnosis Code:** C7A.011 (Malignant carcinoid tumor of the jejunum)
- FL80 Product Information:** LANREOTIDE (lanreotide acetate), 120mg, 1 unit, SQ, NDC 69097-0870-67

FL42 Revenue Code –

List revenue codes in ascending order. Enter the appropriate numeric revenue code in F: 42 to explain each charge in FL 47.²¹

FL43 Revenue Code Description –

Provide a narrative description or standard abbreviation for each revenue code shown in FL 42 on the adjacent line in FL 43. If an NDC is required, submit "N4" followed by the 11-digit NCD, quantity qualifier, and units of measure.²²

FL44 Drug Code –

Enter the appropriate CPT and HCPCS codes and modifiers, as required by the specific payer. **Drug –** J3490 (unclassified drugs) **Administration –** 96402 (Chemotherapy administration, subcutaneous or intramuscular; hormonal anti-neoplastic)²³

FL45 Date of Service –

Enter the date of service for which LANREOTIDE was administered²⁴

FL46 Service Units –

Enter the number of service units for each line item. For a miscellaneous code, bill 1 unit²⁵

FL66 Diagnosis Code –

Enter the appropriate ICD-10-CM diagnosis code(s) for the patient. **For example:** C7A.011 (Malignant carcinoid tumor of the jejunum)²⁶

FL80 Product Information –

When using a miscellaneous code to report the use of LANREOTIDE, include the NDC number (i.e., 69097-0870-67), name of the drug, strength of drug administered, and dose.²⁷

Note: When billing for unlisted drugs using a NOC code, form locator 80 (or the equivalent electronic Loop/Segment in the 837-I, must be populated with the specific name of the drug(s) and/or a short descriptor of the drug(s).²⁸

IMPORTANT SAFETY INFORMATION (continued):

Most Common Adverse Reactions (>10%) include

abdominal pain, musculoskeletal pain, vomiting, headache, injection site reaction, hyperglycemia, hypertension, and cholelithiasis.

Immunogenicity: There is potential for immunogenicity.

Drug Interactions

Insulin and Oral Hypoglycemic Drugs: Blood glucose levels should be monitored and antidiabetic treatment should be adjusted accordingly.

Cyclosporine: Lanreotide Injection may decrease the absorption of cyclosporine. Dosage adjustment of cyclosporine may be needed.

Bromocriptine: Lanreotide may increase the absorption of bromocriptine.

Bradycardia-Inducing Drugs (e.g., beta-blockers):

Lanreotide Injection may decrease heart rate. Dosage adjustment of the coadministered drug may be necessary.

Drug Metabolism Interactions: Avoid other drugs mainly metabolized by CYP3A4 and which have a low therapeutic index (e.g., quinidine, terfenadine). Drugs metabolized by the liver may be metabolized more slowly during Lanreotide Injection treatment; dose reductions of the concomitantly administered medications should be considered.

Use in Specific Populations

Pregnancy: The risk of major birth defects and miscarriage is unknown.

Lactation: Advise women not to breastfeed during treatment with Lanreotide Injection and for 6 months following the last dose.

Females and Males of Reproductive Potential: Lanreotide Injection may reduce fertility in females of reproductive potential.

Geriatric Use: Dose selection for an elderly patient should be cautious.

You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Cipla at 1-866-604-3268.

Please see the full Prescribing Information for additional Important Safety Information.

REFERENCES:

1. <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&AppNo=215395>.
2. LANREOTIDE PRESCRIBING INFORMATION, CIPLA. 2021.
3. Lanreotide Customer Notice 01 Mar 2022, HCPCs Decision Tree, p2.
4. Noridian Healthcare Solutions – Unlisted and Not Otherwise Classified Code Billing/Section Unclassified Drug Billing.
5. Medicare Physician Fee Schedule.
6. Medicare Claims Processing Manual, Chapter 17 – DRUGS AND BIOLOGICALS/Section 20.1.3, p14/¶2.
7. Redbook LANREOTIDE Pricing Search.
8. Medicare Claims Processing Manual, Chapter 17 – DRUGS AND BIOLOGICALS/Section 20.1.3, p13/¶4/lines.
9. Centers for Medicare & Medicaid Services. ICD-10-CM tabular list of diseases and injuries.
10. American Medical Association. CPT® 2021 Standard Edition: eBook (Non-printable). Chicago, IL: American Medical Association; 2021.
11. CMS-1500.
12. CMS Internet Only Manual (IOM), Publication 100-04, Medicare Claims Processing Manual, Chapter 26/ Section 10.4/p12.
13. CMS Internet Only Manual (IOM), Publication 100-04, Medicare Claims Processing Manual, Chapter 26/ Section 10.4/p14.
14. CMS Internet Only Manual (IOM), Publication 100-04, Medicare Claims Processing Manual, Chapter 26/ Section 10.4/p15-16.
15. CMS Internet Only Manual (IOM), Publication 100-04, Medicare Claims Processing Manual, Chapter 26/ Section 10.4/p16.
16. CMS Internet Only Manual (IOM), Publication 100-04, Medicare Claims Processing Manual, Chapter 26/ Section 10.4/p16-17.
17. CMS Internet Only Manual (IOM), Publication 100-04, Medicare Claims Processing Manual, Chapter 26/ Section 10.4/p17.
18. NUCC 1500 – 837P Crosswalk.
19. CMS-1450.
20. Medicare Billing: Form CMS-1450 and the 837 Institutional.
21. Medicare Claims Processing Manual Chapter 25 – Completing and Processing the Forms CMS-1450 Data Set/ Section 75.4/ p15.
22. Medicare Claims Processing Manual Chapter 25 – Completing and Processing the Forms CMS-1450 Data Set/ Section 75.5/ p16.
23. Medicare Claims Processing Manual Chapter 25 – Completing and Processing the Forms CMS-1450 Data Set/ Section 75.5/ p16-17.
24. Medicare Claims Processing Manual Chapter 25 – Completing and Processing the Forms CMS-1450 Data Set/ Section 75.5/ p17-18.
25. Medicare Claims Processing Manual Chapter 25 – Completing and Processing the Forms CMS-1450 Data Set/ Section 75.5/ p18.
26. Medicare Claims Processing Manual Chapter 25 – Completing and Processing the Forms CMS-1450 Data Set/ Section 75.6/ p21.
27. Medicare Claims Processing Manual Chapter 25 – Completing and Processing the Forms CMS-1450 Data Set/ Section 75.6/ p24.
28. Medicare Billing: Form CMS-1450 and the 837 Institutional.
29. https://www.cms.gov/Medicare/Coding/MedHCPCSGenInfo/Downloads/HCPCS_Decision_Tree_and_Definitions.pdf?msclkid=4def1237cfd711ec9994472674288896

[Click here for Full Prescribing Information.](#)

www.ciplalanreotide.com

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LANREOTIDE INJECTION safely and effectively. See full prescribing information for LANREOTIDE INJECTION.

LANREOTIDE injection, for subcutaneous use

Initial U.S. Approval: 2007

INDICATIONS AND USAGE

Lanreotide Injection is a somatostatin analog indicated for:

- The long-term treatment of acromegalic patients who have had an inadequate response to or cannot be treated with surgery and/or radiotherapy. (1.1)
- The treatment of adult patients with unresectable, well- or moderately-differentiated, locally advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NETs) to improve progression-free survival. (1.2)

DOSAGE AND ADMINISTRATION

Administration (2.1):

- For deep subcutaneous injection only.
- Intended for administration by a healthcare provider.
- Administer in the superior external quadrant of the buttock.
- Alternate injection sites.

Recommended Dosage (2.1)

- Acromegaly: 90 mg every 4 weeks for 3 months. Adjust thereafter based on GH and/or IGF-1 levels. See full prescribing information for titration regimen.
- GEP-NETs: 120 mg every 4 weeks.

Dosage Adjustment:

- See full prescribing information for dosage adjustment in patients with acromegaly and renal or hepatic impairment. (2.2, 2.3)

DOSAGE FORMS AND STRENGTHS

Injection: 60 mg/0.2 mL, 90 mg/0.3 mL, and 120 mg/0.5 mL of lanreotide in single-dose prefilled syringes (3)

CONTRAINDICATIONS

Hypersensitivity to lanreotide. (4)

WARNINGS AND PRECAUTIONS

- **Cholelithiasis and Complications of Cholelithiasis:** Monitor periodically. Discontinue if complications of cholelithiasis are suspected. Gallstones may occur; consider periodic monitoring. (5.1)
- **Hyperglycemia and Hypoglycemia:** Glucose monitoring is recommended and antidiabetic treatment adjusted accordingly. (5.2, 7.1)
- **Cardiovascular Abnormalities:** Decrease in heart rate may occur. Use with caution in at-risk patients. (5.3)
- **Thyroid Function Abnormalities:** Decreases in thyroid function may occur; perform tests where clinically indicated. (5.4)

ADVERSE REACTIONS

Most common adverse reactions are:

- **Acromegaly (>5%):** diarrhea, cholelithiasis, abdominal pain, nausea and injection site reactions. (6.1)
- **GEP-NET (>10%):** abdominal pain, musculoskeletal pain, vomiting, headache, injection site reaction, hyperglycemia, hypertension, and cholelithiasis. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Cipla Ltd. Inc. at 1-866-604-3268 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

DRUG INTERACTIONS

- **Cyclosporine:** Lanreotide Injection may decrease the absorption of cyclosporine. Dosage adjustment of cyclosporine may be needed. (7.2)
- **Bromocriptine:** Lanreotide Injection may increase the absorption of bromocriptine. (7.3)
- **Bradycardia-Inducing Drugs (e.g., beta-blockers):** Lanreotide Injection may decrease heart rate. Dosage adjustment of the coadministered drug may be necessary. (7.3)

USE IN SPECIFIC POPULATIONS

Lactation: Advise women not to breastfeed during treatment and for 6 months after the last dose. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 12/2021

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE	7.5	Drug Metabolism Interactions
1.1	Acromegaly	8	USE IN SPECIFIC POPULATIONS
1.2	Gastroenteropancreatic Neuroendocrine Tumors	8.1	Pregnancy
2	DOSAGE AND ADMINISTRATION	8.2	Lactation
2.1	Recommended Dosage	8.3	Females and Males of Reproductive Potential
2.2	Dosage Adjustment in Renal Impairment	8.4	Pediatric Use
2.3	Dosage Adjustment in Hepatic Impairment	8.5	Geriatric Use
2.4	Important Administration Instructions	8.6	Renal Impairment
3	DOSAGE FORMS AND STRENGTHS	8.7	Hepatic Impairment
4	CONTRAINDICATIONS	11	DESCRIPTION
5	WARNINGS AND PRECAUTIONS	12	CLINICAL PHARMACOLOGY
5.1	Cholelithiasis and Complications of Cholelithiasis	12.1	Mechanism of Action
5.2	Hyperglycemia and Hypoglycemia	12.2	Pharmacodynamics
5.3	Cardiovascular Abnormalities	12.3	Pharmacokinetics
5.4	Thyroid Function Abnormalities	13	NONCLINICAL TOXICOLOGY
5.5	Monitoring: Laboratory Tests	13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
6	ADVERSE REACTIONS	14	CLINICAL STUDIES
6.1	Clinical Trials Experience	14.1	Acromegaly
6.2	Immunogenicity	14.2	Gastroenteropancreatic Neuroendocrine Tumors
6.3	Postmarketing Experience	16	HOW SUPPLIED/STORAGE AND HANDLING
7	DRUG INTERACTIONS	17	PATIENT COUNSELING INFORMATION
7.1	Insulin and Oral Hypoglycemic Drugs		
7.2	Cyclosporine		
7.3	Bromocriptine		
7.4	Bradycardia-Inducing Drugs		

* Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Acromegaly

Lanreotide Injection is indicated for the long-term treatment of acromegalic patients who have had an inadequate response to surgery and/or radiotherapy, or for whom surgery and/or radiotherapy is not an option.

The goal of treatment in acromegaly is to reduce growth hormone (GH) and insulin growth factor-1 (IGF-1) levels to normal.

1.2 Gastroenteropancreatic Neuroendocrine Tumors

Lanreotide Injection is indicated for the treatment of adult patients with unresectable, well or moderately differentiated, locally advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NETs) to improve progression-free survival.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

Acromegaly

The recommended starting dosage of Lanreotide Injection is 90 mg given via the deep subcutaneous route, at 4-week intervals for 3 months.

After 3 months, the Lanreotide injection dosage may be adjusted as follows:

- GH greater than 1 ng/mL to less than or equal to 2.5 ng/mL, IGF-1 normal, and clinical symptoms controlled: maintain dosage at 90 mg every 4 weeks.
- GH greater than 2.5 ng/mL, IGF-1 elevated, and/or clinical symptoms uncontrolled: increase dosage to 120 mg every 4 weeks.
- GH less than or equal to 1 ng/mL, IGF-1 normal, and clinical symptoms controlled: reduce dosage to 60 mg every 4 weeks.

Thereafter, the dosage should be adjusted according to the response of the patient as judged by a reduction in serum GH and/or IGF-1 levels; and/or changes in symptoms of acromegaly.

Patients who are controlled on Lanreotide Injection 60 or 90 mg may be considered for an extended dosing interval of Lanreotide Injection 120 mg every 6 or 8 weeks. GH and IGF-1 levels should be obtained 6 weeks after this change in dosing regimen to evaluate persistence of patient response.

Continued monitoring of patient response with dosage adjustments for biochemical and clinical symptom control, as necessary, is recommended.

Gastroenteropancreatic Neuroendocrine Tumors (GEP-NETs)

The recommended dosage of Lanreotide Injection is 120 mg administered every 4 weeks by deep subcutaneous injection.

2.2 Dosage Adjustment in Renal Impairment

Acromegaly

The recommended starting dosage of Lanreotide Injection in acromegalic patients with moderate or severe renal impairment (creatinine clearance less than 60 mL/min) is 60 mg via the deep subcutaneous route at 4-week intervals for 3 months followed by dosage adjustment [see *Dosage and Administration (2.1), Use in Specific Populations (8.6)*].

2.3 Dosage Adjustment in Hepatic Impairment

Acromegaly

The recommended starting dosage of Lanreotide Injection in acromegalic patients with moderate or severe hepatic impairment (Child-Pugh Class B or C) is 60 mg via the deep subcutaneous route at 4-week intervals for 3 months followed by dosage adjustment [see *Dosage and Administration (2.1), Use in Specific Populations (8.7)*].

2.4 Important Administration Instructions

The following instructions explain how to inject Lanreotide Injection:

1. Read all instructions carefully before starting the injection. Follow this procedure exactly, as it may differ from your past experience.
2. This is a single-dose pre-filled syringe with a single-use safety needle with a green needle shield (that cannot be removed) in a clear needle cap.
3. ALL the medication must be injected SLOWLY over 20 seconds during the use.
4. If you drop or damage the device in any way, please call 1-866-604-3268.









Figure 1

The box includes the following items:

- Sterile needle pack containing **Sterile** needle
- **Sterile** Laminated pouch with sterile syringe pre-filled with LANREOTIDE INJECTION
- Instructions for Use Leaflet
- Prescribing Information Leaflet



CAUTION

 <p>DO NOT TOUCH THE GREEN NEEDLE SHIELD. THIS IS NOT A CAP.</p>	<ul style="list-style-type: none">• NEVER TOUCH OR TRY TO OPEN THE GREEN NEEDLE SHIELD throughout the course of operation of the device.• Green needle shield is NOT a removable cap or cover for the needle.• Green needle shield will automatically activate once the injection is complete.• Green needle shield is a self-operating safety lock mechanism.• Needle is fully covered by green needle shield.• Needle is visible only through a small window in the green needle shield.
 <p>Plunger</p> <p>Syringe Barrel</p> <p>20 sec</p>	<ul style="list-style-type: none">• Inject medication slowly over 20 seconds.• DO NOT rush the injection.
 <p>30:00 min.</p>	<ul style="list-style-type: none">• Remove box from refrigerator 30 minutes prior to injecting.• Product left in its sealed pouch at room temperature (not to exceed 104°F or 40°C) for up to 24 hours may be returned to the refrigerator for continued storage and usage at a later time.
 <p>OK</p>	<ul style="list-style-type: none">• Stretch out the skin around injection site to make it flat and tight using your thumb and index finger.• DO NOT pinch the skin around injection site.

A. Before you start




Figure 2



Figure 3




Figure 4


A1. Remove **LANREOTIDE INJECTION** from the refrigerator 30 minutes prior to injecting (**Figure 2**). Do **not** open the sterile pouch yet.

Note: Product left in its sealed pouch at room temperature (not to exceed 104° F or 40 °C) for up to 24 hours may be returned to the refrigerator for continued storage and use at a later time.

A2. Wash your hands with soap and dry your hands thoroughly before starting (**Figure 3**).

Follow the doctor or institution's policy on the use of surgical gloves during the procedure

A3. Before opening the sterile pouch, confirm that it is intact, and that the medication has not expired. **The Syringe is sterile only if the pouch is sealed and undamaged. Do not use if the pouch is opened, tampered with or damaged.** The expiry date is printed on the outer carton and the pouch - **Do not use if the medication has expired.** (**Figure 4**).



Or



Figure 5




Figure 6



Figure 7

A4. Make sure there is a clean surface for preparation.

Find a clean, comfortable area for the patient to relax during procedure (**Figure 5**).

It's important that the patient remains as still as possible during the injection.

A5. Choose injection site - the sites are upper outer areas of the **buttock** as shown below
It is very important that you only inject in one of the areas marked OK in the picture (**Figure 6**).

Alternate the injection site between the right and left side each time an injection of **LANREOTIDE INJECTION** is administered.

A7. Tear open the sterile pouch and take out the sterile pre-filled syringe (**Figure 8**).




Figure 8

B. Prepare the syringe

B1: Open the sterile needle cap (Figure 9)

- The needle is sterile only if the needle cap is sealed and undamaged. Do not use if the needle cap is opened, tampered with or damaged.
- Hold the clear needle cap and pull the lid off.
- **DO NOT TOUCH THE OPEN END OF THE NEEDLE CAP TO MAINTAIN STERILITY.**




Figure 9

B2: Remove the cap from the sterile syringe (Figure 10)

- With one hand, hold the syringe barrel steady (**not the plunger**).
- With the other hand, remove the cap by twisting it.




Figure 10

B3: Prepare the assembly (Figure 11)

- Hold the needle cap with one hand and the syringe barrel (**not the plunger**) with the other
- Carefully insert the open end of the syringe into the open end of the needle cap
- Twist the syringe barrel clockwise until it is tight to make sure that the syringe is well connected to the safety needle.
- **ENSURE THAT BOTH PARTS OF THE DEVICE (SYRINGE AND NEEDLE) ARE COMPLETELY CONNECTED.**
The assembly is fully locked when you cannot turn it any further.




Figure 11


B4: Remove the needle from the needle cap

- Hold the syringe barrel (**not the plunger**).
- Pull the needle straight out from the needle cap **without twisting or turning** (Figure 12).



Figure 12

CAUTION: NEVER TOUCH THE GREEN NEEDLE SHIELD. THERE IS A RISK OF NEEDLE STICK INJURY. (Figure 13)



- Green needle shield is a self-operating safety lock mechanism and is not a removable cover or cap for the needle. It will activate once the injection is complete.
- The needle is fully covered by the green shield and is visible only through the small round window at the end of the shield.

Figure 13

C. Perform injection

C1: Position the syringe assembly



Figure 14

- Stretch out the skin around the injection site to make it flat and tight using your thumb and index finger. (Figure 14)



Figure 15

DO NOT pinch the skin (Figure 15)




Figure 16



Figure 17

C2: Insert the needle (Figure 18)




Figure 18

- Holding the skin flat and tight, push the syringe assembly firmly into the skin.
- Do NOT pinch the skin.
- **Keep pushing until you reach a “hard stop” and only the green collar at the end of the green needle shield remains visible.**
- **Do NOT** push the plunger yet.
- **DO NOT aspirate.**
- Keep pressing against the skin.
- During this step you should not see the needle.

C3: Push the top of the plunger (Figure 19)



Figure 19

- Keep holding your hand on the syringe barrel to maintain a 90-degree angle throughout the injection.
- While keeping the syringe pressed against the skin, **slowly** press down the plunger. You may want to use both hands while applying pressure during the injection of drug.
- Continue slowly pushing the plunger over 20 seconds until the Plunger top touches the syringe end .
- **DO NOT** rush the injection. The medication is thicker and harder to push than you might expect. Pushing the plunger too fast may cause discomfort to the patient.

- While depressing the plunger, slowly count to 20 and **CONTINUE STEADY PRESSURE** on the plunger. You may find it helpful to say:
 - “1 one-thousand”
 - “2 one-thousand”
 - “3 one-thousand” up to “20 one-thousand”
 - During this step, as the needle is lowering, the green needle shield will retract. You should only see the green collar of the green needle shield.
 - During this step you should not see the needle.

D. Remove and throw away the syringe assembly




Figure 20

D1: Remove from the skin (Figure 20)

- Lift the syringe straight up and away from the body.
- **The green needle shield will return to its original position and will fully cover the needle.**



Figure 21

D2: Apply gentle pressure (Figure 21)

- Apply gentle pressure to the injection site with a dry cotton ball or sterile gauze to prevent any bleeding.
- **Do NOT** rub or massage the injection site after administration.




Figure 22

D3: Check needle (Figure 22)

- Check through the green needle shield’s round windows that the needle was not damaged during administration.
- If any damage or malfunction to the needle is suspected please contact 1-866-604-3268



Figure 23

D4: Throw away / Disposal (Figure 23)

- DO NOT remove needle from syringe
- Dispose of complete product in sharps disposal container
- Do not dispose of the syringe or needle in the regular trash.
- The syringe and needle are not reusable.

Issued: 12/2021
SAP Code for IFU - 99325175

3 DOSAGE FORMS AND STRENGTHS

Injection: 60 mg/0.2 mL, 90 mg/0.3 mL, and 120 mg/0.5 mL of lanreotide provided as lanreotide acetate, in single-dose, prefilled syringes packaged with a safety needle. The prefilled syringes contain a white to pale yellow, semi-solid formulation.

4 CONTRAINDICATIONS

Lanreotide Injection is contraindicated in patients with history of a hypersensitivity to lanreotide. Allergic reactions (including angioedema and anaphylaxis) have been reported following administration of lanreotide [*see Adverse Reactions (6.3)*].

5 WARNINGS AND PRECAUTIONS

5.1 Cholelithiasis and Complications of Cholelithiasis

Lanreotide may reduce gallbladder motility and lead to gallstone formation; therefore, patients may need to be monitored periodically [*see Adverse Reactions (6.1), Clinical Pharmacology (12.2)*]. There have been postmarketing reports of cholelithiasis (gallstones) resulting in complications, including cholecystitis, cholangitis, and pancreatitis, and requiring cholecystectomy in patients taking lanreotide. If complications of cholelithiasis are suspected, discontinue Lanreotide Injection and treat appropriately.

5.2 Hyperglycemia and Hypoglycemia

Pharmacological studies in animals and humans show that lanreotide, like somatostatin and other somatostatin analogs, inhibits the secretion of insulin and glucagon. Hence, patients treated with Lanreotide Injection may experience hypoglycemia or hyperglycemia. Blood glucose levels should be monitored when Lanreotide Injection treatment is initiated, or when the dose is altered, and antidiabetic treatment should be adjusted accordingly [*see Adverse Reactions (6.1)*].

5.3 Cardiovascular Abnormalities

The most common overall cardiac adverse reactions observed in three pooled lanreotide cardiac studies in patients with acromegaly were sinus bradycardia (12/217, 5.5%),

bradycardia (6/217, 2.8%), and hypertension (12/217, 5.5%) [*see Adverse Reactions (6.1)*].

In 81 patients with baseline heart rates of 60 beats per minute (bpm) or greater treated with lanreotide in Study 3, the incidence of heart rate less than 60 bpm was 23% (19/81) as compared to 16% (15/94) of placebo treated patients; 10 patients (12%) had documented heart rates less than 60 bpm on more than one visit. The incidence of documented episodes of heart rate less than 50 bpm as well as the incidence of bradycardia reported as an adverse event was 1% in each treatment group. Initiate appropriate medical management in patients who develop symptomatic bradycardia.

In patients without underlying cardiac disease, lanreotide may lead to a decrease in heart rate without necessarily reaching the threshold of bradycardia. In patients suffering from cardiac disorders prior to Lanreotide Injection treatment, sinus bradycardia may occur. Care should be taken when initiating treatment with Lanreotide Injection in patients with bradycardia.

5.4 Thyroid Function Abnormalities

Slight decreases in thyroid function have been seen during treatment with lanreotide in acromegalic patients, though clinical hypothyroidism is rare (less than 1%). Thyroid function tests are recommended where clinically indicated.

5.5 Monitoring: Laboratory Tests

Acromegaly: Serum GH and IGF-1 levels are useful markers of the disease and the effectiveness of treatment [*see Dosage and Administration (2.1)*].

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Cholelithiasis and Complications of Cholelithiasis [*see Warnings and Precautions (5.1)*]
- Hyperglycemia and Hypoglycemia [*see Warnings and Precautions (5.2)*]
- Cardiovascular Abnormalities [*see Warnings and Precautions (5.3)*]
- Thyroid Function Abnormalities [*see Warnings and Precautions (5.4)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Acromegaly

The data described below reflect exposure to lanreotide in 416 acromegalic patients in seven studies. One study was a fixed-dose pharmacokinetic study. The other six studies were open-label or extension studies, one had a placebo-controlled, run-in period, and another had an active control. The population was mainly Caucasian (329/353, 93%) with a median age of 53 years of age (range 19 to 84 years). Fifty-four subjects (13%) were age 66 to 74 and 18 subjects (4.3%) were 75 years of age and older.

Patients were evenly matched for sex (205 males and 211 females). The median average monthly dose was 91.2 mg (e.g., 90 mg injected via the deep subcutaneous route every 4 weeks) over 385 days with a median cumulative dose of 1290 mg. Of the patients reporting acromegaly, severity at baseline (N=265), serum GH levels were less than 10 ng/mL for 69% (183/265) of the patients and 10 ng/mL or greater for 31% (82/265) of the patients.

The most commonly reported adverse reactions reported by greater than 5% of patients who received lanreotide (N=416) in the overall pooled safety studies in acromegaly patients were gastrointestinal disorders (diarrhea, abdominal pain, nausea, constipation, flatulence, vomiting, loose stools), cholelithiasis, and injection site reactions.

Tables 1 and 2 present adverse reaction data from clinical studies with lanreotide in acromegalic patients. The tables include data from a single clinical study and pooled data from seven clinical studies.

Adverse Reactions in Parallel Fixed-Dose Phase of Study 1

The incidence of treatment-emergent adverse reactions for lanreotide 60, 90, and 120 mg by dose as reported during the first 4 months (fixed-dose phase) of Study 1 [see *Clinical Studies (14.1)*] are provided in Table 1.

Table 1: Adverse Reactions at an Incidence of Greater than 5% with Lanreotide Overall and Occurring at Higher Rate than Placebo: Placebo-Controlled and Fixed-Dose Phase of Study 1 By Dose

Body System Preferred Term	Placebo-Controlled Double-Blind Phase Weeks 0 to 4		Fixed-Dose Phase Double-Blind + Single-Blind Weeks 0 to 20			
	Placebo (N=25)	Lanreotide Overall (N=83)	Lanreotide 60 mg (N=34)	Lanreotide 90 mg (N=36)	Lanreotide 120 mg (N=37)	Lanreotide Overall (N=107)
	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
Gastrointestinal System Disorders	1 (4%)	30 (36%)	12 (35%)	21 (58%)	27 (73%)	60 (56%)
Diarrhea	0	26 (31%)	9 (26%)	15 (42%)	24 (65%)	48 (45%)
Abdominal pain	1 (4%)	6 (7%)	3 (9%)	6 (17%)	7 (19%)	16 (15%)
Flatulence	0	5 (6%)	0 (0%)	3 (8%)	5 (14%)	8 (7%)
Application Site Disorders (Injection site mass/ pain/ reaction/ inflammation)	0 (0%)	5 (6%)	3 (9%)	4 (11%)	8 (22%)	15 (14%)
Liver and Biliary System Disorders	1 (4%)	3 (4%)	9 (26%)	7 (19%)	4 (11%)	20 (19%)
Cholelithiasis	0	2 (2%)	5 (15%)	6 (17%)	3 (8%)	14 (13%)
Heart Rate & Rhythm Disorders	0	8 (10%)	7 (21%)	2 (6%)	5 (14%)	14 (13%)
Bradycardia	0	7 (8%)	6 (18%)	2 (6%)	2 (5%)	10 (9%)
Red Blood Cell Disorders	0	6 (7%)	2 (6%)	5 (14%)	2 (5%)	9 (8%)
Anemia	0	6 (7%)	2 (6%)	5 (14%)	2 (5%)	9 (8%)
Metabolic & Nutritional Disorders	3 (12%)	13 (16%)	8 (24%)	9 (25%)	4 (11%)	21 (20%)
Weight decrease	0	7 (8%)	3 (9%)	4 (11%)	2 (5%)	9 (8%)

A patient is counted only once for each body system and preferred term.
Dictionary = WHOART.

In Study 1, the adverse reactions of diarrhea, abdominal pain, and flatulence increased in incidence with increasing dose of lanreotide injection .

Adverse Reactions in Long-Term Clinical Trials

Table 2 provides the most common adverse reactions (greater than 5%) that occurred in 416 acromegalic patients treated with lanreotide pooled from 7 studies compared to those patients from the 2 efficacy studies (Studies 1 and 2). Patients with elevated GH and IGF-1 levels were either naive to somatostatin analog therapy or had undergone a 3-month washout [see *Clinical Studies (14.1)*].

Table 2: Adverse Reactions in Lanreotide -Treated Patients at an Incidence Greater than 5% in Overall Group Versus Adverse Reactions Reported in Studies 1 and 2

System Organ Class	Number and Percentage of Patients			
	Studies 1 & 2 (N=170)		Overall Pooled Data (N=416)	
	N	%	N	%
Patients with any Adverse Reactions	157	92	356	86
Gastrointestinal disorders	121	71	235	57
Diarrhea	81	48	155	37
Abdominal pain	34	20	79	19
Nausea	15	9	46	11
Constipation	9	5	33	8
Flatulence	12	7	30	7
Vomiting	8	5	28	7
Loose stools	16	9	23	6
Hepatobiliary disorders	53	31	99	24
Cholelithiasis	45	27	85	20
General disorders and administration site conditions	51	30	91	22
(Injection site pain /mass /induration/ nodule/pruritus)	28	17	37	9
Musculoskeletal and connective tissue disorders	44	26	70	17
Arthralgia	17	10	30	7
Nervous system disorders	34	20	80	19
Headache	9	5	30	7

Dictionary = MedDRA 7.1

In addition to the adverse reactions listed in Table 2, the following reactions were also seen:

- Sinus bradycardia occurred in 7% (12) of patients in the pooled Study 1 and 2 and in 3% (13) of patients in the overall pooled studies.
- Hypertension occurred in 7% (11) of patients in the pooled Study 1 and 2 and in 5% (20) of patients in the overall pooled studies.
- Anemia occurred in 7% (12) of patients in the pooled Study 1 and 2 and in 3% (14) of patients in the overall pooled studies.

Gastrointestinal Adverse Reactions

In the pooled clinical studies of lanreotide therapy, a variety of gastrointestinal (GI) reactions occurred, the majority of which were mild to moderate in severity. One percent of acromegalic patients treated with lanreotide in the pooled clinical studies discontinued treatment because of gastrointestinal reactions.

Pancreatitis was reported in less than 1% of patients.

Gallbladder Adverse Reactions

In clinical studies involving 416 acromegalic patients treated with lanreotide, cholelithiasis and gallbladder sludge were reported in 20% of the patients. Among 167 acromegalic patients treated with lanreotide who underwent routine evaluation with gallbladder ultrasound, 17% had gallstones at baseline. New cholelithiasis was reported in 12% of patients. Cholelithiasis may be related to dose or duration of exposure [see *Warnings and Precautions (5.1)*].

Injection Site Reactions

In the pooled clinical studies, injection site pain (4%) and injection site mass (2%) were the

most frequently reported local adverse drug reactions that occurred with the administration of lanreotide. In a specific analysis, 20 of 413 patients (5%) presented indurations at the injection site. Injection site adverse reactions were more commonly reported soon after the start of treatment and were less commonly reported as treatment continued. Such adverse reactions were usually mild or moderate but did lead to withdrawal from clinical studies in two subjects.

Glucose Metabolism Adverse Reactions

In the clinical studies in acromegalic patients treated with lanreotide, adverse reactions of dysglycemia (hypoglycemia, hyperglycemia, diabetes) were reported by 14% (47/332) of patients and were considered related to study drug in 7% (24/332) of patients [see *Warnings and Precautions (5.2)*].

Cardiac Adverse Reactions

In the pooled clinical studies, sinus bradycardia (3%) was the most frequently observed heart rate and rhythm disorder. All other cardiac adverse drug reactions were observed in less than 1% of patients. The relationship of these events to lanreotide could not be established because many of these patients had underlying cardiac disease [see *Warnings and Precautions (5.3)*].

A comparative echocardiography study of lanreotide and another somatostatin analog demonstrated no difference in the development of new or worsening valvular regurgitation between the 2 treatments over 1 year. The occurrence of clinically significant mitral regurgitation (i.e., moderate or severe in intensity) or of clinically significant aortic regurgitation (i.e., at least mild in intensity) was low in both groups of patients throughout the study.

Other Adverse Reactions

For the most commonly occurring adverse reactions in the pooled analysis, diarrhea, abdominal pain, and cholelithiasis, there was no apparent trend for increasing incidence with age. GI disorders and renal and urinary disorders were more common in patients with documented hepatic impairment; however, the incidence of cholelithiasis was similar between groups.

Gastroenteropancreatic Neuroendocrine Tumors

The safety of lanreotide 120 mg for the treatment of patients with gastroenteropancreatic neuroendocrine tumors (GEP-NETs) was evaluated in Study 3, a double-blind, placebo-controlled trial. Patients in Study 3 were randomized to receive lanreotide (N=101) or placebo (N=103) administered by deep subcutaneous injection once every 4 weeks. The data below reflect exposure to lanreotide in 101 patients with GEP-NETs, including 87 patients exposed for at least 6 months and 72 patients exposed for at least 1 year (median duration of exposure 22 months). Patients treated with lanreotide had a median age of 64 years (range 30 to 83 years), 53% were men and 96% were Caucasian. Eighty-one percent of patients (83/101) in the lanreotide arm and 82% of patients (82/103) in the placebo arm did not have disease progression within 6 months of enrollment and had not received prior therapy for GEP-NETs. The rates of discontinuation due to treatment-emergent adverse reactions were 5% (5/101 patients) in the lanreotide arm and 3% (3/103 patients) in the placebo arm.

Table 3 compares the adverse reactions reported with an incidence of 5% and greater in patients receiving lanreotide 120 mg administered every 4 weeks and reported more commonly than placebo.

Table 3: Adverse Reactions Occurring in 5% and Greater of Lanreotide - Treated Patients and at a Higher Rate Than in Placebo -Treated Patients in Study 3

Adverse Reaction	Lanreotide 120 mg N=101		Placebo N=103	
	Any (%)	Severe** (%)	Any (%)	Severe** (%)
Any Adverse Reactions	88	26	90	31
Abdominal pain ¹	34*	6*	24*	4
Musculoskeletal pain ²	19*	2*	13	2
Vomiting	19*	2*	9*	2*
Headache	16	0	11	1
Injection site reaction ³	15	0	7	0
Hyperglycemia ⁴	14*	0	5	0
Hypertension ⁵	14*	1*	5	0
Cholelithiasis	14*	1*	7	0
Dizziness	9	0	2*	0
Depression ⁶	7	0	1	0
Dyspnea	6	0	1	0

¹ Includes preferred terms of abdominal pain, abdominal pain upper/lower, abdominal discomfort

² Includes preferred terms of myalgia, musculoskeletal discomfort, musculoskeletal pain, back pain

³ Includes preferred terms of infusion site extravasation, injection site discomfort, injection site granuloma, injections site hematoma, injection site hemorrhage, injection site induration, injection site mass, injections site nodule, injection site pain, injection site pruritus, injection site rash, injection site reaction, injection site swelling

⁴ Includes preferred terms of diabetes mellitus, glucose tolerance impaired, hyperglycemia, type 2 diabetes mellitus

⁵ Includes preferred terms of hypertension, hypertensive crisis

⁶ Includes preferred terms of depression, depressed mood

* Includes one or more serious adverse events (SAEs) defined as any event that results in death, is life threatening, results in hospitalization or prolongation of hospitalization, results in persistent or significant disability, results in congenital anomaly/birth defect, or may jeopardize the patient and may require medical or surgical intervention to prevent one of the outcomes listed.** Defined as hazardous to well-being, significant impairment of function or incapacitation

6.2 Immunogenicity

As with all peptides, there is potential for immunogenicity. The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies in the studies described below with the incidence of antibodies in other studies or to other lanreotide products may be misleading.

Laboratory investigations of acromegalic patients treated with lanreotide in clinical studies show that the percentage of patients with putative antibodies at any time point after treatment is low (less than 1% to 4% of patients in specific studies whose antibodies were tested). The antibodies did not appear to affect the efficacy or safety of lanreotide.

In Study 3, development of anti-lanreotide antibodies was assessed using a radioimmuno-precipitation assay. In patients with GEP NETs receiving lanreotide, the incidence of anti-lanreotide antibodies was 4% (3 of 82) at 24 weeks, 10% (7 of 67) at 48 weeks, 11% (6 of 57) at 72 weeks, and 10% (8 of 84) at 96 weeks. Assessment for neutralizing antibodies was not conducted.

6.3 Post-marketing Experience

The following adverse reactions have been identified during post-approval use of lanreotide. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Hepatobiliary: steatorrhea; cholecystitis, cholangitis, pancreatitis, which have sometimes required cholecystectomy

Hypersensitivity: angioedema and anaphylaxis

Injection site reactions: injection site abscess

7 DRUG INTERACTIONS

7.1 Insulin and Oral Hypoglycemic Drugs

Lanreotide, like somatostatin and other somatostatin analogs, inhibits the secretion of insulin and glucagon. Therefore, blood glucose levels should be monitored when Lanreotide Injection treatment is initiated or when the dose is altered, and antidiabetic treatment should be adjusted accordingly [see *Warnings and Precautions (5.2)*].

7.2 Cyclosporine

Concomitant administration of cyclosporine with Lanreotide Injection may decrease the absorption of cyclosporine, and therefore, may necessitate adjustment of cyclosporine dose to maintain therapeutic drug concentrations. [see *Clinical Pharmacology (12.3)*]

7.3 Bromocriptine

Limited published data indicate that concomitant administration of a somatostatin analog and bromocriptine may increase the absorption of bromocriptine [see *Clinical Pharmacology (12.3)*].

7.4 Bradycardia-Inducing Drugs

Concomitant administration of bradycardia-inducing drugs (e.g., beta-blockers) may have an additive effect on the reduction of heart rate associated with lanreotide. Dosage adjustments of concomitant drugs may be necessary.

7.5 Drug Metabolism Interactions

The limited published data available indicate that somatostatin analogs may decrease the metabolic clearance of compounds known to be metabolized by cytochrome P450 enzymes, which may be due to the suppression of growth hormone. Since it cannot be excluded that Lanreotide Injection may have this effect, avoid other drugs mainly metabolized by CYP3A4 and which have a low therapeutic index (e.g., quinidine, terfenadine). Drugs metabolized by the liver may be metabolized more slowly during Lanreotide Injection treatment and dose reductions of the concomitantly administered medications should be considered [see *Clinical Pharmacology (12.3)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Limited available data based on post-marketing case reports with lanreotide use in pregnant women are not sufficient to determine a drug-associated risk of adverse developmental outcomes. In animal reproduction studies, decreased embryo/fetal survival was observed in pregnant rats and rabbits at subcutaneous doses 5- and 2-times the maximum recommended human dose (MRHD) of 120 mg, respectively (*see Data*).

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Animal Data

A reproductive study in pregnant rats given 30 mg/kg of lanreotide by subcutaneous injection every 2 weeks (5 times the human dose, based on body surface area comparisons) resulted in decreased embryo/fetal survival. A study in pregnant rabbits given subcutaneous injections of 0.45 mg/kg/day (2 times the human therapeutic exposures at the maximum recommended dose of 120 mg, based on comparisons of relative body surface area) shows decreased fetal survival and increased fetal skeletal/soft tissue abnormalities.

8.2 Lactation

Risk Summary

There is no information available on the presence of lanreotide in human milk, the effects of the drug on the breastfed infant, or the effects of the drug on milk production. Studies show that lanreotide administered subcutaneously passes into the milk of lactating rats; however, due to species-specific differences in lactation physiology, animal data may not reliably predict drug levels in human milk. Because of the potential for serious adverse reactions in breastfed infants from Lanreotide Injection, including effects on glucose metabolism and bradycardia, advise women not to breastfeed during treatment with Lanreotide Injection and for 6 months (6 half-lives) following the last dose.

8.3 Females and Males of Reproductive Potential

Infertility

Females

Based on results from animal studies conducted in female rats, Lanreotide Injection may reduce fertility in females of reproductive potential [*see Nonclinical Toxicology (13.1)*].

8.4 Pediatric Use

The safety and effectiveness of Lanreotide Injection in pediatric patients have not been established.

8.5 Geriatric Use

No overall differences in safety or effectiveness were observed between elderly patients with acromegaly compared with younger patients and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. Studies 3 and 4, conducted in patients with neuroendocrine tumors, did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients.

Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.6 Renal Impairment

Acromegaly

Lanreotide has been studied in patients with end-stage renal function on dialysis, but has not been studied in patients with mild, moderate, or severe renal impairment. It is recommended that patients with moderate or severe renal impairment receive a starting dose of Lanreotide Injection of 60 mg. Caution should be exercised when considering patients with moderate or severe renal impairment for an extended dosing interval of Lanreotide Injection 120 mg every 6 or 8 weeks [see *Dosage and Administration (2.4) and Clinical Pharmacology (12.3)*].

Neuroendocrine Tumors (NET) – Gastroenteropancreatic Neuroendocrine Tumors

No effect was observed in total clearance of lanreotide in patients with mild to moderate renal impairment receiving lanreotide 120 mg. Patients with severe renal impairment were not studied [see *Clinical Pharmacology (12.3)*].

8.7 Hepatic Impairment

Acromegaly

It is recommended that patients with moderate or severe hepatic impairment receive a starting dose of Lanreotide Injection 60 mg. Caution should be exercised when considering patients with moderate or severe hepatic impairment for an extended dosing interval of Lanreotide Injection 120 mg every 6 or 8 weeks [see *Dosage and Administration (2.4) and Clinical Pharmacology (12.3)*].

Neuroendocrine Tumors (NET) – Gastroenteropancreatic Neuroendocrine Tumors

Lanreotide has not been studied in patients with hepatic impairment.


11 DESCRIPTION

Lanreotide Injection 60 mg/0.2 mL, 90 mg/0.3 mL, and 120 mg/0.5 mL is a prolonged-release formulation for deep subcutaneous injection. It contains the drug substance lanreotide acetate, a synthetic octapeptide with a biological activity similar to naturally occurring somatostatin, water for injection and acetic acid (for pH adjustment).

Lanreotide Injection is available as sterile, ready-to-use, single-dose prefilled syringes containing lanreotide acetate supersaturated bulk solution of 24.6% w/w lanreotide base.

Each syringe contains:	Lanreotide Injection 60 mg/0.2 mL	Lanreotide Injection 90 mg/0.3 mL	Lanreotide Injection 120 mg/0.5 mL
Lanreotide acetate	89.9 mg	123.2 mg	156.6 mg
Acetic Acid	q.s.	q.s.	q.s.
Water for injection	236.4 mg	324.1 mg	411.6 mg
Total Weight	328.9 mg	450.9 mg	572.8 mg

Lanreotide acetate is a synthetic cyclical octapeptide analog of the natural hormone, somatostatin. Lanreotide acetate is chemically known as [cyclo S-S]-3-(2-naphthyl)-D-alanyl-L-cysteinyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-cysteinyl-L-threoninamide, acetate salt. Its molecular weight is 1096.34 (base) and its amino acid sequence is:



The Lanreotide Injection in the prefilled syringe is a white to pale yellow, semi-solid formulation.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Lanreotide, the active component of Lanreotide Injection is an octapeptide analog of natural somatostatin. The mechanism of action of lanreotide is believed to be similar to that of natural somatostatin.

12.2 Pharmacodynamics

Lanreotide has a high affinity for human somatostatin receptors (SSTR) 2 and 5 and a reduced binding affinity for human SSTR1, 3, and 4. Activity at human SSTR2 and 5 is the primary mechanism believed responsible for GH inhibition. Like somatostatin, lanreotide is an inhibitor of various endocrine, neuroendocrine, exocrine, and paracrine functions.

The primary pharmacodynamic effect of lanreotide is a reduction of GH and/or IGF-1 levels enabling normalization of levels in acromegalic patients [see *Clinical Studies (14.1)*]. In acromegalic patients, lanreotide reduces GH levels in a dose-dependent way. After a single injection of lanreotide, plasma GH levels fall rapidly and are maintained for at least 28 days.

Lanreotide inhibits the basal secretion of motilin, gastric inhibitory peptide, and pancreatic polypeptide, but has no significant effect on the secretion of secretin. Lanreotide inhibits postprandial secretion of pancreatic polypeptide, gastrin, and cholecystokinin (CCK). In healthy subjects, lanreotide produces a reduction and a delay in postprandial insulin secretion, resulting in transient, mild glucose intolerance.

Lanreotide inhibits meal-stimulated pancreatic secretions, and reduces duodenal bicarbonate and amylase concentrations, and produces a transient reduction in gastric acidity.

Lanreotide has been shown to inhibit gallbladder contractility and bile secretion in healthy subjects [see *Warnings and Precautions (5.1)*].

In healthy subjects, lanreotide inhibits meal-induced increases in superior mesenteric artery and portal venous blood flow but has no effect on basal or meal-stimulated renal blood flow. Lanreotide has no effect on renal plasma flow or renal vascular resistance. However, a

transient decrease in glomerular filtration rate (GFR) and filtration fraction has been observed after a single injection of lanreotide.

In healthy subjects, non-significant reductions in glucagon levels were seen after lanreotide administration. In diabetic non-acromegalic subjects receiving a continuous infusion (21-day) of lanreotide, serum glucose concentrations were temporarily decreased by 20% to 30% after the start and end of the infusion. Serum glucose concentrations returned to normal levels within 24 hours. A significant decrease in insulin concentrations was recorded between baseline and Day 1 only [see *Warnings and Precautions (5.2)*].

Lanreotide inhibits the nocturnal increase in thyroid-stimulating hormone (TSH) seen in healthy subjects. Lanreotide reduces prolactin levels in acromegalic patients treated on a long-term basis [see *Warnings and Precautions (5.4)*].

12.3 Pharmacokinetics

Lanreotide Injection is thought to form a drug depot at the injection site due to the interaction of the formulation with physiological fluids. The most likely mechanism of drug release is a passive diffusion of the precipitated drug from the depot towards the surrounding tissues, followed by the absorption to the bloodstream.

After a single, deep subcutaneous administration, the mean absolute bioavailability of lanreotide in healthy subjects was 73.4, 69.0, and 78.4% for the 60 mg, 90 mg, and 120 mg doses, respectively. Mean C_{max} values ranged from 4.3 to 8.4 ng/mL during the first day. Single-dose linearity was demonstrated with respect to AUC and C_{max} and showed high inter-subject variability. Lanreotide showed sustained release of lanreotide with a half-life of 23 to 30 days. Mean serum concentrations were > 1 ng/mL throughout 28 days at 90 mg and 120 mg and > 0.9 ng/mL at 60 mg.

In studies evaluating excretion, <5% of lanreotide was excreted in urine and less than 0.5% was recovered unchanged in feces, indicative of some biliary excretion.

Acromegaly

In a repeat-dose administration pharmacokinetics (PK) study in acromegalic patients, rapid initial release was seen giving peak levels during the first day after administration. At doses of lanreotide between 60 and 120 mg, linear pharmacokinetics were observed in acromegalic patients. At steady state, mean C_{max} values were 3.8 ± 0.5 , 5.7 ± 1.7 , and 7.7 ± 2.5 ng/mL, increasing linearly with dose. The mean accumulation ratio index was 2.7, which is in line with the range of values for the half-life of lanreotide. The steady-state trough serum lanreotide concentrations in patients receiving lanreotide every 28 days were 1.8 ± 0.3 ; 2.5 ± 0.9 and 3.8 ± 1.0 ng/mL at 60 mg, 90 mg, and 120 mg doses, respectively. A limited initial burst effect and a low peak-to-trough fluctuation (81% to 108%) of the serum concentration at the plateau were observed.

For the same doses, similar values were obtained in clinical studies after at least four administrations (2.3 ± 0.9 , 3.2 ± 1.1 , and 4.0 ± 1.4 ng/mL, respectively).

Pharmacokinetic data from studies evaluating extended dosing use of lanreotide 120 mg demonstrated mean steady-state, C_{min} values between 1.6 and 2.3 ng/mL for the 8- and 6-week treatment interval, respectively.

Gastroenteropancreatic Neuroendocrine Tumors

In patients with GEP-NETs treated with lanreotide 120 mg every 4 weeks, steady state concentrations were reached after 4 to 5 injections and the mean trough serum lanreotide concentrations at steady state ranged from 5.3 to 8.6 ng/mL.

Specific Populations

Lanreotide has not been studied in specific populations. However, the pharmacokinetics of lanreotide in renal impaired, hepatic impaired, and geriatric subjects were evaluated after IV administration of lanreotide immediate release formulation (IRF) at 7 mcg/kg dose.

Geriatric

Studies in healthy elderly subjects showed an 85% increase in half-life and a 65% increase in mean residence time (MRT) of lanreotide compared to those seen in healthy young subjects; however, there was no change in either AUC or C_{max} of lanreotide in elderly as compared to healthy young subjects. Age has no effect on clearance of lanreotide based on population PK analysis in patients with GEP-NET which included 122 patients aged 65 to 85 years with neuroendocrine tumors.

Renal Impairment

An approximate 2-fold decrease in total serum clearance of lanreotide, with a consequent 2-fold increase in half-life and AUC was observed [see *Dosage and Administration (2.2) and Use in Specific Populations (8.6)*].

Mild (CL_{cr} 60-89 mL/min) or moderate (CL_{cr} 30-59 mL/min) renal impairment has no effect on clearance of lanreotide in patients with GEP-NET based on population PK analysis which included 106 patients with mild and 59 patients with moderate renal impairment treated with lanreotide. GEP-NET patients with severe renal impairment (CL_{cr} < 30 mL/min) were not studied.

Hepatic Impairment

In subjects with moderate to severe hepatic impairment, a 30% reduction in clearance of lanreotide was observed [see *Dosage and Administration (2.3) and Use in Specific Populations (8.7)*].

The effect of hepatic impairment on clearance of lanreotide has not been studied in patients with GEP-NET.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Standard lifetime carcinogenicity bioassays were conducted in mice and rats. Mice were given daily subcutaneous doses of lanreotide at 0.5, 1.5, 5, 10, and 30 mg/kg for 104 weeks. Cutaneous and subcutaneous tumors of fibrous connective tissues at the injection sites were observed at the high dose of 30 mg/kg/day. Fibrosarcomas in both genders and malignant fibrous histiocytomas were observed in males at 30 mg/kg/day resulting in exposures 3 times higher than the clinical therapeutic exposure at the maximum therapeutic dose of 120 mg given by monthly subcutaneous injection based on the AUC values. Rats were given daily subcutaneous doses of lanreotide at 0.1, 0.2, and 0.5 mg/kg for 104 weeks. Increased cutaneous and subcutaneous tumors of fibrous connective tissues at the injection sites were observed at the dose of 0.5 mg/kg/day resulting in exposures less than the clinical therapeutic exposure at 120 mg given by monthly subcutaneous injection. The increased incidence of injection site tumors in rodents is likely related to the increased dosing frequency (daily) in animals compared to monthly dosing in humans and therefore may not be clinically relevant.

Lanreotide was not genotoxic in tests for gene mutations in a bacterial mutagenicity (Ames) assay, or mouse lymphoma cell assay with or without metabolic activation. Lanreotide was not genotoxic in tests for the detection of chromosomal aberrations in a human lymphocyte and *in vivo* mouse micronucleus assay.

In a fertility study conducted with lanreotide in rats, reduced female fecundity was observed at estimated exposure corresponding to approximately 10-fold the plasma exposure at the MRHD of 120 mg. The fertility of male rats was unaffected by the treatment up to an estimated exposure corresponding to approximately 11-fold the plasma exposure at the MRHD of 120 mg.

14 CLINICAL STUDIES

14.1 Acromegaly

The effect of lanreotide on reducing GH and IGF-levels and control of symptoms in patients with acromegaly was studied in 2 long-term, multiple-dose, randomized, multicenter studies.

Study 1

This 1-year study included a 4-week, double-blind, placebo-controlled phase; a 16-week single-blind, fixed-dose phase; and a 32-week, open-label, dose-titration phase. Patients with active acromegaly, based on biochemical tests and medical history, entered a 12-week washout period if there was previous treatment with a somatostatin analog or a dopaminergic agonist.

Upon entry, patients were randomly allocated to receive a single, deep subcutaneous injection of lanreotide 60, 90, or 120 mg or placebo. Four weeks later, patients entered a fixed-dose phase where they received 4 injections of lanreotide followed by a dose-titration phase of 8 injections for a total of 13 injections over 52 weeks (including the placebo phase). Injections were given at 4-week intervals. During the dose-titration phase of the study, the dose was titrated twice (every fourth injection), as needed, according to individual GH and IGF-1 levels.

A total of 108 patients (51 males, 57 females) were enrolled in the initial placebo-controlled phase of the study. Half (54/108) of the patients had never been treated with a somatostatin analog or dopamine agonist or had stopped treatment for at least 3 months prior to their participation in the study, and were required to have a mean GH level greater than 5 ng/mL at their first visit. The other half of the patients had received prior treatment with a somatostatin analog or a dopamine agonist before study entry and at study entry were required to have a mean GH concentration greater than 3 ng/mL and at least a 100% increase in mean GH concentration after washout of medication.

One hundred and seven (107) patients completed the placebo-controlled phase, 105 patients completed the fixed-dose phase, and 99 patients completed the dose-titration phase. Patients not completing withdrew due to adverse events (5) or lack of efficacy (4).

In the double-blind phase of Study 1, a total of 52 (63%) of the 83 lanreotide-treated patients had a greater than 50% decrease in mean GH from baseline to Week 4, including 52%, 44%, and 90% of patients in the 60, 90, and 120 mg groups, respectively, compared to placebo (0%, 0/25). In the fixed-dose phase at Week 16, 72% of all 107 lanreotide-treated patients had a decrease from baseline in mean GH of greater than 50%, including 68% (23/34), 64% (23/36), and 84% (31/37) of patients in the 60, 90, and 120 mg lanreotide treatment groups, respectively. Efficacy achieved in the first 16 weeks was maintained for the duration of the study (see Table 4).

Table 4: Overall Efficacy Results Based on GH and IGF-1 Levels by Treatment Phase in Study 1

		Baseline N=107	Before Titration 1 (16 weeks) N=107	Before Titration 2 (32 weeks) N=105	Last Value Available* N=107
GH					
≤5.0 ng/mL	Number of Responders (%)	20 (19%)	72 (67%)	76 (72%)	74 (69%)
≤2.5 ng/mL	Number of Responders (%)	0 (0%)	52 (49%)	59 (56%)	55 (51%)
≤1.0 ng/mL	Number of Responders (%)	0 (0%)	15 (14%)	18 (17%)	17 (16%)
Median GH	ng/mL	10.27	2.53	2.20	2.43
GH Reduction	Median % Reduction	--	75.5	78.2	75.5
IGF-1					
Normal³	Number of Responders (%)	9 (8%)	58 (54%)	57 (54%)	62 (58%)
Median IGF-1	ng/mL	775.0	332.0 ¹	316.5 ²	326.0
IGF-1 Reduction	Median % Reduction	--	52.3 ¹	54.5 ²	55.4
IGF-1 Normal³ + GH ≤2.5 ng/mL	Number of Responders (%)	0 (0%)	41 (38%)	46 (44%)	44 (41%)

¹ n=105, ² n=102, ³ Age-adjusted

*Last Observation Carried Forward

Study 2

This was a 48-week, open-label, uncontrolled, multicenter study that enrolled patients who had an IGF-1 concentration 1.3 times or greater than the upper limit of the normal age-adjusted range. Patients receiving treatment with a somatostatin analog (other than lanreotide) or a dopaminergic agonist had to attain this IGF-1 concentration after a washout period of up to 3 months.

Patients were initially enrolled in a 4-month, fixed-dose phase where they received 4 deep subcutaneous injections of lanreotide 90 mg, at 4-week intervals. Patients then entered a dose-titration phase where the dose of lanreotide was adjusted based on GH and IGF-1 levels at the beginning of the dose-titration phase and, if necessary, again after another 4 injections. Patients titrated up to the maximum dose (120 mg) were not allowed to titrate down again.

A total of 63 patients (38 males, 25 females) entered the fixed-dose phase of the trial and 57 patients completed 48 weeks of treatment. Six patients withdrew due to adverse reactions (3), other reasons (2), or lack of efficacy (1).

After 48 weeks of treatment with lanreotide at 4-week intervals, 43% (27/63) of the acromegalic patients in this study achieved normal age-adjusted IGF-1 concentrations. Mean IGF-1 concentrations after treatment completion were 1.3 ± 0.7 times the upper limit of normal compared to 2.5 ± 1.1 times the upper limit of normal at baseline.

The reduction in IGF-1 concentrations over time correlated with a corresponding marked decrease in mean GH concentrations. The proportion of patients with mean GH concentrations less than 2.5 ng/mL increased significantly from 35% to 77% after the fixed-dose phase and 85% at the end of the study. At the end of treatment, 24/63 (38%) of patients had both normal IGF-1 concentrations and a GH concentration of less than or equal to 2.5 ng/mL (see Table 5) and 17/63 patients (27%) had both normal IGF-1 concentrations and a GH concentration of less than 1 ng/mL.

Table 5: Overall Efficacy Results Based on GH and IGF-1 Levels by Treatment Phase in Study 2

		Baseline N=63	Before Titration 1 (12 wks) N=63	Before Titration 2 (28 wks) N=59	Last Value Available* N=63
IGF-1					
Normal¹	Number of Responders (%)	0 (0%)	17 (27%)	22 (37%)	27 (43%)
Median IGF-1	ng/mL	689.0	382.0	334.0	317.0
IGF-1 Reduction	Median % Reduction	--	41.0	51.0	50.3
GH					
≤5.0 ng/mL	Number of Responders (%)	40 (64%)	59 (94%)	57 (97%)	62 (98%)
≤2.5 ng/mL	Number of Responders (%)	21 (33%)	47 (75%)	47 (80%)	54 (86%)
≤1.0 ng/mL	Number of Responders (%)	8 (13%)	19 (30%)	18 (31%)	28 (44%)
Median GH	ng/ml	3.71	1.65	1.48	1.13
GH Reduction	Median % Reduction	--	63.2	66.7	78.6 ²
IGF-1 normal¹ + GH ≤2.5 ng/mL	Number of Responders (%)	0 (0%)	14 (22%)	20 (34%)	24 (38%)

¹ Age-adjusted, ²N= 62,

* Last Observation Carried Forward

Examination of age and gender subgroups did not identify differences in response to lanreotide among these subgroups. The limited number of patients in the different racial subgroups did not raise any concerns regarding efficacy of lanreotide in these subgroups.

14.2 Gastroenteropancreatic Neuroendocrine Tumors

The efficacy of lanreotide was established in a multicenter, randomized, double-blind, placebo-controlled trial of 204 patients with unresectable, well or moderately differentiated, metastatic or locally advanced, gastroenteropancreatic neuroendocrine tumors. Patients were required to have non-functioning tumors without hormone-related symptoms. Patients were randomized 1:1 to receive lanreotide 120 mg (n=101) or placebo (n=103) every 4 weeks until disease progression, unacceptable toxicity, or a maximum of 96 weeks of treatment. Randomization was stratified by the presence or absence of prior therapy and by the presence or absence of disease progression within 6 months of enrollment. The major efficacy outcome measure was progression-free survival (PFS), defined as time to disease progression as assessed by central independent radiological review using the Response Evaluation Criteria in Solid Tumors (RECIST 1.0) or death.

The median patient age was 63 years (range 30 to 92 years) and 95% were Caucasian. Disease progression was present in nine of 204 patients (4.4%) in the 6 months prior to enrollment and 29 patients (14%) received prior chemotherapy. Ninety-one patients (45%) had primary sites of disease in the pancreas, with the remainder originating in the midgut (35%), hindgut (7%), or unknown primary location (13%). The majority (69%) of the study population had grade 1 tumors. Baseline prognostic characteristics were similar between arms

with one exception: there were 39% of patients in the lanreotide arm and 27% of patients in the placebo arm who had hepatic involvement by tumor of greater than 25%.

Patients in the lanreotide arm had a statistically significant improvement in PFS compared to patients receiving placebo (see Table 6 and Figure 1).


Table 6: Efficacy Results in Study 3

	Lanreotide n=101	Placebo n=103
Number of Events (%)	32 (31.7%)	60 (58.3%)
Median PFS (months)(95% CI)	NE ¹ (NE, NE)	16.6 (11.2, 22.1)
HR (95% CI)	0.47 (0.30, 0.73) ²	
Log-rank p-value	<0.001	

¹ NE = not reached at 22 months

² Hazard Ratio is derived from a Cox stratified proportional hazards model

Figure 1: Kaplan-Meier Curves of Progression-Free Survival



16 HOW SUPPLIED/STORAGE AND HANDLING

Lanreotide Injection is supplied in strengths of 60 mg/0.2 mL, 90 mg/0.3 mL, and 120 mg/0.5 mL as a white to pale yellow, semi-solid formulation in a single-dose, sterile, prefilled, ready-to-use, polypropylene syringe and a safety needle device. The safety needle device is a sterile, single use needle system consisting of a needle (1.2 mm x 20 mm, stainless steel) held in protective plastic safety housing.

The single-dose prefilled syringe is contained in a plastic tray and is packed in a triple-layered aluminium pouch. The sterile safety needle is co-packaged along with the sealed aluminium pouch in the kit carton box and is attached to the former at the point of use.

NDC 69097-880-67
NDC 69097-890-67
NDC 69097-870-67

60 mg/0.2 mL, sterile, prefilled syringe
90 mg/0.3 mL, sterile, prefilled syringe
120 mg/0.5 mL, sterile, prefilled syringe

Storage and Handling

Store Lanreotide Injection in the refrigerator at 2°C to 8°C (36°F to 46°F). Protect from light. Store in the original package.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Hypersensitivity Reactions

Advise patients to immediately contact their healthcare provider if they experience serious hypersensitivity reactions, such as angioedema or anaphylaxis [*see Contraindications (4)*].

Cholelithiasis and Complications of Cholelithiasis

Advise patients to contact their healthcare provider if they experience signs or symptoms of gallstones (cholelithiasis) or complications of gallstones (e.g., cholecystitis, cholangitis, or pancreatitis) [*see Warnings and Precautions (5.1)*].

Hyperglycemia and Hypoglycemia

Advise patients to immediately contact their healthcare provider if they experience signs or symptoms of hyper- or hypoglycemia [*see Warnings and Precautions (5.2)*].

Cardiovascular Abnormalities

Advise patients to immediately contact their healthcare provider if they experience bradycardia [*see Warnings and Precautions (5.3)*].

Thyroid Function Abnormalities

Advise patients to contact their healthcare provider if they experience signs or symptoms of hypothyroidism [*see Warnings and Precautions (5.4)*].

Laboratory Tests

Advise patients with acromegaly that response to Lanreotide Injection should be monitored by periodic measurements of GH and IGF-1 levels, with a goal of decreasing these levels to the normal range [*see Dosage and Administration (2.1)*].

Lactation

Advise women not to breastfeed during treatment with Lanreotide Injection and for 6 months after the last dose [*see Use in Specific Populations (8.2)*].

Infertility

Advise females of reproductive potential of the potential for reduced fertility from Lanreotide Injection [*see Use in Specific Populations (8.3)*].

Manufactured by:
Pharmathen International S.A.
Industrial Park Sapas,
Rodopi Prefecture, Block No 5,
Rodopi 69300,
Greece

Manufactured for:
Cipla USA Inc.
10 Independence Boulevard, Suite 300
Warren, NJ 07059

PATIENT INFORMATION
Lanreotide (lan-REE-oh-tide) Injection
for subcutaneous use

Read this Patient Information before you receive your first Lanreotide injection and before each injection. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is Lanreotide injection?

Lanreotide injection is a prescription medicine used for:

- the long-term treatment of people with acromegaly when:
 - surgery or radiotherapy have not worked well enough or
 - they are not able to have surgery or radiotherapy.
- the treatment of adults with a type of cancer known as neuroendocrine tumors, from the gastrointestinal tract or the pancreas (GEP-NETs) that has spread or cannot be removed by surgery.

It is not known if Lanreotide injection is safe and effective in children.

Who should not receive Lanreotide injection?

Do not receive Lanreotide injection if you are allergic to lanreotide.

What should I tell my healthcare provider before receiving Lanreotide injection?

Before you receive Lanreotide injection, tell your healthcare provider about all of your medical conditions, including if you:

- have gallbladder problems.
- have diabetes.
- have heart problems.
- have thyroid problems.
- have kidney problems.
- have liver problems.
- are pregnant or plan to become pregnant. It is not known if Lanreotide injection will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Lanreotide injection passes into your breast milk. You should not breastfeed if you receive Lanreotide injection and for 6 months after your last dose of Lanreotide injection.
- are a person who can become pregnant. Lanreotide injection may affect fertility and may affect your ability to become pregnant. Talk to your healthcare provider if this is a concern for you.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Lanreotide injection and other medicines may affect each other, causing side effects. Lanreotide injection may affect the way other medicines work, and other medicines may affect how Lanreotide injection works. Your dose of Lanreotide injection or your other medicines may need to be changed.

Especially tell your healthcare provider if you take:

- insulin or other diabetes medicines.
- cyclosporine (Gengraf, Neoral, or Sandimmune).
- medicines that lower your heart rate such as beta blockers.

How will I receive Lanreotide injection?

- You will receive a Lanreotide injection dose every 4 weeks in your healthcare provider's office.
- Your healthcare provider may change your dose of Lanreotide injection or the length of time between your injections. Your healthcare provider will tell you how long you need to receive Lanreotide injection.
- Lanreotide injection is injected deep under the skin of the upper outer area of your buttock. Your injection site should change (alternate) between your right and left buttock from one injection of Lanreotide injection to the next.
- During your treatment with Lanreotide injection for acromegaly, your healthcare provider may do certain blood tests to see if Lanreotide injection is working.

What should I avoid while receiving Lanreotide injection?

Lanreotide injection can cause dizziness. If you have dizziness, do not drive a car or operate machinery.

What are the possible side effects of Lanreotide injection? Lanreotide injection

may cause serious side effects, including:

- **Gallstones (cholelithiasis) and complications that can happen if you have gallstones.** Gallstones are a serious but common side effect in people who take Lanreotide injection and have acromegaly and GEP-NET. Your healthcare provider may check your gallbladder before and during treatment with Lanreotide injection. Possible complications of gallstones include inflammation and infection of the gallbladder and pancreatitis. Tell your healthcare provider if you get any symptoms of gallstones, including:
 - sudden pain in your upper right stomach area (abdomen)
 - sudden pain in your right shoulder or between your shoulder blades
 - yellowing of your skin and whites of your eyes
 - fever with chills

- nausea

- **Changes in your blood sugar** (high blood sugar or low blood sugar). If you have diabetes, test your blood sugar as your healthcare provider tells you to. Your healthcare provider may change your dose of diabetes medicine especially when you first start receiving Lanreotide injection or if your dose of Lanreotide injection changes. High blood sugar is a common side effect in people with GEP-NET.

Tell your healthcare provider right away if you have any signs or symptoms of high blood sugar or low blood sugar.

Signs and symptoms of high blood sugar may include:

- increased thirst
- increased appetite
- nausea
- weakness or tiredness
- urinating more often than normal
- your breath smells like fruit

Signs and symptoms of low blood sugar may include:

- dizziness or lightheadedness
- sweating
- confusion
- headache
- blurred vision
- slurred speech
- shakiness
- fast heartbeat
- irritability or mood changes
- hunger

- **Slow heart rate.** Tell your healthcare provider right away if you have slowing of your heart rate or if you have symptoms of a slow heart rate, including:

- dizziness or lightheadedness
- fainting or near-fainting
- chest pain
- shortness of breath
- confusion or memory problems
- weakness, extreme tiredness

- **High blood pressure.** High blood pressure can happen in people who receive Lanreotide injection and is a common side effect in people with GEP-NET.

- **Changes in thyroid function.** Lanreotide injection can cause the thyroid gland to not make enough thyroid hormones that the body needs (hypothyroidism) in people who have acromegaly. Tell your healthcare provider if you have signs and symptoms of low thyroid hormones levels, including:

- fatigue
- weight gain
- a puffy face
- being cold all of the time
- constipation
- dry skin
- thinning, dry hair
- decreased sweating
- depression

The most common side effects of Lanreotide injection in people with acromegaly include:

- diarrhea
- stomach area (abdominal) pain
- nausea
- pain, itching, or a lump at the injection site

The most common side effects of Lanreotide injection in people with GEP-NET include:

- stomach area (abdominal) pain
- muscle and joint aches
- vomiting
- headache
- pain, itching, or a lump at the injection site

Tell your healthcare provider right away if you have signs of an allergic reaction after receiving Lanreotide injection, including:

- swelling of your face, lips, mouth or tongue
- breathing problems
- fainting, dizziness, feeling lightheaded (low blood pressure)
- itching
- flushing or redness of your skin
- rash
- hives

These are not all the possible side effects of Lanreotide injection. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of Lanreotide injection.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not receive Lanreotide injection for a condition for which it was not prescribed. You can ask your healthcare provider for information about Lanreotide injection that is written for health professionals.

What are the ingredients in Lanreotide injection?

Active ingredient: lanreotide acetate

Inactive ingredients: water for injection and acetic acid (for pH adjustment)

Manufactured by: Pharmathen International S.A., Rodopi, Greece

Manufactured for: Cipla USA, Inc., 10 Independence Boulevard, Suite 300, Warren, NJ 07059

SAP Code: 99332893

For more information, go to www.ciplausa.com or call 1-886-604-3268.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Issued: 12/2021